

CONVERGENCE OF A SINGULAR EULER-POISSON APPROXIMATION OF THE INCOMPRESSIBLE NAVIER-STOKES EQUATIONS

ROBERTO NATALINI[†] AND FRÉDÉRIC ROUSSET[‡]

ABSTRACT. In this note, we rigorously justify a singular approximation of the incompressible Navier-Stokes equations. Our approximation combines two classical approximations of the incompressible Euler equations: a standard relaxation approximation, but with a diffusive scaling, and the Euler-Poisson equations in the quasineutral regime.

1. INTRODUCTION

We shall consider the following system:

$$(1) \quad \begin{cases} \partial_t u^\varepsilon + \operatorname{div} V^\varepsilon = \nabla \varphi^\varepsilon, \\ \varepsilon \partial_t V^\varepsilon + a^2 \nabla u^\varepsilon = u^\varepsilon \otimes u^\varepsilon - V^\varepsilon, \\ \partial_t \rho^\varepsilon + \operatorname{div} (\rho^\varepsilon u^\varepsilon) = 0, \\ \Delta \varphi^\varepsilon = \frac{\rho^\varepsilon - 1}{\varepsilon}, \end{cases}$$

for $t > 0$, $x \in \mathbb{T}^d$, $u^\varepsilon \in \mathbb{R}^d$, $V^\varepsilon \in \mathbb{R}^{d,d}$, $\rho^\varepsilon \in \mathbb{R}$. Here $\varepsilon > 0$ is small parameter and a is a non zero constant value. To solve uniquely the Poisson equation, we add the condition $\int_{\mathbb{T}^d} \rho = 1$. Passing to the limit when ε goes to zero, it is easy to see, at least at a very formal level, that $(\rho^\varepsilon, u^\varepsilon, V^\varepsilon)$ tends to $(\rho^{NS}, u^{NS}, V^{NS})$, where

$$(2) \quad \rho^{NS} = 1, \quad V^{NS} = u^{NS} \otimes u^{NS} - a^2 \nabla u^{NS}$$

and

$$(3) \quad \begin{cases} \partial_t u^{NS} + \operatorname{div} (u^{NS} \otimes u^{NS}) = a^2 \Delta u^{NS} + \nabla \varphi^{NS}, \\ \operatorname{div} u^{NS} = 0. \end{cases}$$

In other words, u^{NS} is a solution of the incompressible Navier-Stokes equations. The aim of this note is to give a rigorous justification to this formal computation. We shall prove the following result.

Theorem 1. *Let u^{NS} a solution of the Navier-Stokes equations (3) such that $u^{NS} \in C([0, T], H^{m+3}(\mathbb{T}^d))$ and $\int_{\mathbb{T}^d} u^{NS} = 0$ for $d \leq 3$ and $m > 1 + d/2$. Assume that the*

1991 *Mathematics Subject Classification.* Primary: 35Q30; Secondary: 76D05.

Key words and phrases. Incompressible Navier-Stokes equations, quasineutral regime, Euler-Poisson equations, diffusive relaxation approximations, hyperbolic singular perturbations.

The research activity reported in this paper has been partially conducted within the European Union RTN HYKE project: HPRN-CT-2002-00282.

[†] Istituto per le Applicazioni del Calcolo “Mauro Picone”, Consiglio Nazionale delle Ricerche, Viale del Policlinico, 137, I-00161 Roma, Italy, r.natalini@iac.cnr.it

[‡] CNRS, Laboratoire J.-A. Dieudonné, UMR 6621, Université de Nice, Parc Valrose F-06108 Nice Cedex 02, France, frousset@math.unice.fr.

initial value $(u_0^\varepsilon, V_0^\varepsilon, \rho_0^\varepsilon) \in H^{m+1}$ of (1) is such that $\int_{\mathbb{T}^d} \rho_0^\varepsilon = 1$, $\int_{\mathbb{T}^d} u_0^\varepsilon = 0$ and

$$\alpha_m(\varepsilon) := \|u_0^\varepsilon - u_{/t=0}^{NS}\|_{H^{m+1}}^2 + \frac{1}{\varepsilon} \|\rho_0^\varepsilon - 1\|_{H^m}^2 + \varepsilon \|V_0^\varepsilon - V_{/t=0}^{NS}\|_{H^{m+1}}^2 \rightarrow 0$$

when ε goes to zero. Then there exist ε_0 and C_T such that for $0 < \varepsilon \leq \varepsilon_0$ there is a strong solution $(\rho^\varepsilon, u^\varepsilon, V^\varepsilon) \in C([0, T], H^{m+1})$ of (1) such that

$$(4) \quad \|u^\varepsilon(t) - u^{NS}(t)\|_{H^{m+1}}^2 + \frac{1}{\varepsilon} \|\rho^\varepsilon(t) - 1\|_{H^m}^2 \leq C_T(\varepsilon^2 + \alpha_m(\varepsilon)), \quad \forall t \in [0, T].$$

System (1) combines the features of the diffusive relaxation approximation of parabolic systems and of the quasineutral limit of the Euler-Poisson equation towards the incompressible Euler equation. These two problems were much studied by various methods.

Let us recall that the diffusive scaling $(\frac{x}{\sqrt{\varepsilon}}, \frac{t}{\varepsilon})$ has been largely investigated in the framework of hydrodynamic limits of the Boltzmann equations [4] and in the analysis of hyperbolic-parabolic relaxation limits for weak solutions of hyperbolic systems of balance laws with strongly diffusive source terms by means of compensated compactness techniques by Marcatti and collaborators [15, 16, 9]. Under the same scaling, general (possibly degenerate) parabolic equations in multi-D have been approximated by semilinear hyperbolic equations, see [5, 2].

Concerning the quasineutral limit, which consists in the limit of vanishing viscosity for the Poisson part of the system, there are results for various specific models. In particular, this limit has been performed for the Vlasov-Poisson system by Brenier [6], Grenier [11, 12], and in the one dimensional and isothermal Euler-Poisson system by Cordier and Grenier [8]. We refer to [14], [18] and references therein for more recent contributions.

The main focus in the present note is on the use of hyperbolic energy methods for studying incompressible fluids. In this regard, our system can be also seen as a refinement of the relaxation system studied in [7], whose techniques however were restricted to the two dimensional case. In particular our result gives a rate of convergence in strong H^s norm of the solution of the singular system towards a strong solution of the incompressible Navier-Stokes equation. Hyperbolic ideas have been also used in the weak solutions framework by Donatelli and Marcatti to deal with the diffusive relaxation of hyperbolic approximations towards Leray solutions of the incompressible Navier-Stokes equation [10]. Finally, similar approximations have been considered, as reduced kinetic models, in [3] and on a rigorous basis in [13], to design effective numerical schemes for incompressible fluids.

2. PROOF OF THEOREM 1

First, let us set

$$u = u^\varepsilon - u^{NS}, \quad V = V^\varepsilon - V^{NS}, \quad \varphi = \varphi^\varepsilon - \varphi^{NS}, \quad \rho = \rho^\varepsilon - 1.$$

Since the pressure φ^{NS} in the incompressible Navier-Stokes equation is given by

$$\Delta \varphi^{NS} = \nabla(u^{NS} \cdot \nabla u^{NS}) = \nabla u^{NS} \cdot \nabla u^{NS},$$

the vector (ρ, u, V, φ) solves the system

$$(5) \quad \begin{cases} \partial_t u + \operatorname{div} V = \nabla \varphi, \\ \varepsilon \partial_t V + a^2 \nabla u = u^{NS} \otimes u + u \otimes u^{NS} - V + u \otimes u + \varepsilon \partial_t V^{NS}, \\ \partial_t \rho + \operatorname{div} u + u^{NS} \cdot \nabla \rho + \operatorname{div}(\rho u) = 0, \\ \Delta \varphi = \frac{\rho}{\varepsilon} - \nabla u^{NS} \cdot \nabla u^{NS}. \end{cases}$$

As in [14], we make the following change of unknowns:

$$\begin{aligned} d &= \operatorname{div} u, \quad \omega = \operatorname{curl} u, \\ D_i &= (\operatorname{div} V)_i = \sum_j \partial_j V_{ij}, \quad \Omega_{ij} = (\operatorname{curl} V)_{ij} = \partial_i V_j - \partial_j V_i. \end{aligned}$$

By using the last equation in (5) and the fact that $\Delta u = \nabla d - \operatorname{curl} \omega$, we get the following system:

$$\left\{ \begin{aligned} \partial_t d + \operatorname{div} D &= \frac{\rho}{\varepsilon} - \nabla u^{NS} \cdot \nabla u^{NS}, \\ \partial_t \omega + \operatorname{curl} D &= 0, \\ \varepsilon \partial_t D + a^2 (\nabla d - \operatorname{curl} \omega) &= -D + \operatorname{div} (u^{NS} \otimes u + u \otimes u^{NS}) \\ &\quad - \operatorname{div} (u \otimes u) + \varepsilon \partial_t (\operatorname{div} V^{NS}), \\ \varepsilon \partial_t \Omega &= -\Omega + \operatorname{curl} (u^{NS} \otimes u + u \otimes u^{NS}) \\ &\quad + \varepsilon \partial_t \operatorname{curl} V^{NS} - \operatorname{curl} (u \otimes u), \\ \partial_t \rho + u^{NS} \cdot \nabla \rho &= -d - \operatorname{div} (\rho u). \end{aligned} \right.$$

This last system can be written as a singular perturbation of a quasilinear symmetrizable hyperbolic system. Setting $\mathcal{U} = (d, \omega, D, \Omega, \rho)$ yields

$$(6) \quad \mathcal{A}_0^\varepsilon \partial_t \mathcal{U} + \mathcal{A}(t, x, \partial_x) \mathcal{U} + \frac{1}{\varepsilon} \mathcal{K}^\varepsilon \mathcal{U} + \mathcal{D} \mathcal{U} + \mathcal{N}(\mathcal{U}, \partial_x \mathcal{U}) = \mathcal{L}(t, x, \partial_x) \mathcal{U} + \varepsilon R$$

where

$$\begin{aligned} \mathcal{A}_0^\varepsilon &= \operatorname{diag}(1, 1, \varepsilon, \varepsilon, 1), \quad \mathcal{D} = \operatorname{diag}(0, 0, 1, 1, 0), \\ \mathcal{A}(t, x, \partial_x) &= \begin{pmatrix} 0 & 0 & \operatorname{div} & 0 & 0 \\ 0 & 0 & \operatorname{curl} & 0 & 0 \\ a^2 \nabla & -a^2 \operatorname{curl} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & u^{NS} \cdot \nabla \end{pmatrix}, \\ \mathcal{K}^\varepsilon &= \begin{pmatrix} 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \varepsilon & 0 & 0 & 0 & 0 \end{pmatrix}, \\ \mathcal{L}(t, x, \partial_x) \mathcal{U} &= \begin{pmatrix} 0 \\ 0 \\ \operatorname{div} (u^{NS} \otimes u + u \otimes u^{NS}) \\ \operatorname{curl} (u^{NS} \otimes u + u \otimes u^{NS}) \\ 0 \end{pmatrix}, \\ \mathcal{N}(\mathcal{U}, \partial_x \mathcal{U}) &= \begin{pmatrix} 0 \\ 0 \\ \operatorname{div} (u \otimes u) \\ \operatorname{curl} (u \otimes u) \\ \operatorname{div} (\rho u) \end{pmatrix}, \quad R = \begin{pmatrix} -\nabla u^{NS} \cdot \nabla u^{NS} \\ 0 \\ \partial_t \operatorname{div} (V^{NS}) \\ \partial_t \operatorname{curl} (V^{NS}) \\ 0 \end{pmatrix}. \end{aligned}$$

Let us remark that \mathcal{L} is a zero order operator when it acts on \mathcal{U} . The nonlinear term \mathcal{N} can be splitted in

$$\mathcal{N}(\mathcal{U}, \partial_x \mathcal{U}) = \mathcal{S}(\mathcal{U}) + \mathcal{Q}(\mathcal{U}, \partial_x \mathcal{U})$$

where

$$\mathcal{S}(\mathcal{U}) = \begin{pmatrix} 0 \\ 0 \\ \operatorname{div} (u \otimes u) \\ \operatorname{curl} (u \otimes u) \\ \rho \operatorname{div} u \end{pmatrix}$$

is a semilinear term (still considered as acting on \mathcal{U}) and $\mathcal{Q}(\mathcal{U}, \partial_x)\mathcal{U}$ is a quasilinear term with

$$\mathcal{Q}(\mathcal{U}, \partial_x)\mathcal{U} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ u \cdot \nabla \rho \end{pmatrix}.$$

Let us set

$$\Sigma^\varepsilon = \operatorname{diag} (a^2, a^2 I_d, I_d, I_{d \times d}, a^2/\varepsilon)$$

and for $|\alpha| \leq m$

(7)

$$\begin{aligned} E_{\alpha, m}^\varepsilon(t) &= \frac{1}{2} \left(\Sigma^\varepsilon \mathcal{A}_0^\varepsilon \partial^\alpha \mathcal{U}, \partial^\alpha \mathcal{U} \right) \\ &= \frac{1}{2} \left(a^2 \|\partial^\alpha d\|^2 + a^2 \|\partial^\alpha \omega\|^2 + \varepsilon \|\partial^\alpha D\|^2 + \varepsilon \|\partial^\alpha \Omega\|^2 + \frac{a^2}{\varepsilon} \|\partial^\alpha \rho\|^2 \right), \end{aligned}$$

(8)

$$E_m^\varepsilon(t) = \sum_{|\alpha| \leq m} E_{\alpha, m}^\varepsilon.$$

Note that since $\Sigma^\varepsilon \mathcal{Q}(\mathcal{U}, \xi)$ and $\Sigma^\varepsilon \mathcal{A}(t, x, \xi)$ are symmetric, we can consider (6) as a quasilinear hyperbolic system with zero order terms (a linear one coming from \mathcal{L} and a semilinear one coming from \mathcal{S} which are pseudo-differential operators of order zero when acting on \mathcal{U}). Consequently, for $\varepsilon > 0$ fixed, we have a result of local existence and uniqueness of strong solutions in $C_T(H^m)$, see for instance [1, 17]. This allows us to define T^ε as the largest time such that

(9)

$$E_m^\varepsilon(t) \leq M_\varepsilon, \forall t \in [0, T^\varepsilon]$$

where M_ε which is such that $M_\varepsilon \rightarrow 0$ when ε goes to zero will be chosen carefully later. To achieve the proof of Theorem 1, and in particular inequality (4), it is sufficient to establish that $T^\varepsilon \geq T$, which will be proved by showing that in (9) the equality cannot be reached for $T^\varepsilon < T$ thanks to a good choice of M_ε .

Before performing the energy estimate, we apply the operator ∂^α for $|\alpha| \leq m$ to (6), to obtain

(10)

$$\begin{aligned} \mathcal{A}_0^\varepsilon \partial_t \partial^\alpha \mathcal{U} + \mathcal{A}(t, x, \partial_x) \partial^\alpha \mathcal{U} &+ \frac{1}{\varepsilon} \mathcal{K}^\varepsilon \partial^\alpha \mathcal{U} + \mathcal{D} \partial^\alpha \mathcal{U} + \partial^\alpha \mathcal{N}(\mathcal{U}) \\ &= \partial^\alpha \mathcal{L}(t, x, \partial_x) \mathcal{U} + \varepsilon R + [\partial^\alpha, \mathcal{A}(t, x, \partial_x)] \mathcal{U}. \end{aligned}$$

Now, we proceed to perform the energy estimates for (10) in a classical way by taking the scalar product of system (10) with $\Sigma^\varepsilon \partial^\alpha \mathcal{U}$. Along the proof, we shall denote by C a number independent of ε , which actually may change from line to line, and by $\mathcal{C}(\cdot)$ a nondecreasing function. Moreover (\cdot, \cdot) and $\|\cdot\|$ stand for the usual L^2 scalar product and norm, $\|\cdot\|_m$ is the usual H^m Sobolev norm and $\|\cdot\|_{m, \infty}$ is the usual $W^{m, \infty}$ norm.

Let us start the estimate of each term. First, since $\Sigma^\varepsilon \mathcal{A}(x, \xi)$ is symmetric and $\operatorname{div} u^{NS} = 0$, we have that

(11)

$$\left| \left(\Sigma^\varepsilon \mathcal{A}(x, \partial_x) \partial^\alpha \mathcal{U}, \partial^\alpha \mathcal{U} \right) \right| = \left| \frac{a^2}{\varepsilon} \int \operatorname{div} (u^{NS}) |\partial^\alpha \rho|^2 dx \right| = 0.$$

Next, since $\Sigma^\varepsilon \mathcal{K}^\varepsilon$ is skew-symmetric, we have that

$$(12) \quad \left(\Sigma^\varepsilon \mathcal{K}^\varepsilon \partial^\alpha \mathcal{U}, \partial^\alpha \mathcal{U} \right) = 0.$$

Also, we have

$$(13) \quad \left(\Sigma^\varepsilon \mathcal{D} \partial^\alpha \mathcal{U}, \partial^\alpha \mathcal{U} \right) = \|\partial^\alpha D\|^2 + \|\partial^\alpha \Omega\|^2.$$

Now we observe that, since $\int_{\mathbb{T}^d} u = 0$, we have the inequality

$$\|u\|_{m+1}^2 \leq C(\|d\|_m^2 + \|\omega\|_m^2) \leq C\|\nabla u\|_m^2.$$

Hence, the Young inequality gives

$$(14) \quad \begin{aligned} \left(\Sigma^\varepsilon \partial^\alpha \mathcal{L}, \partial^\alpha \mathcal{U} \right) &= \int \partial^\alpha \operatorname{div} (u^{NS} \otimes u + u \otimes u^{NS}) \partial^\alpha D \\ &\quad + \int \partial^\alpha \operatorname{curl} (u^{NS} \otimes u + u \otimes u^{NS}) \partial^\alpha \Omega \\ &\leq \frac{1}{4} (\|\partial^\alpha D\|^2 + \|\partial^\alpha \Omega\|^2) + C(\|u^{NS}\|_{m+1, \infty}) (\|\omega\|_m^2 + \|d\|_m^2) \\ &\leq \frac{1}{4} (\|\partial^\alpha D\|^2 + \|\partial^\alpha \Omega\|^2) + C(\|u^{NS}\|_{m+1, \infty}) E_m^\varepsilon. \end{aligned}$$

To give the estimate of the term $(\Sigma^\varepsilon \partial^\alpha \mathcal{N}, \partial^\alpha \mathcal{U})$, we split it in several terms. The first one is given by, for $m > d/2$,

$$(15) \quad \begin{aligned} &\int \left(\partial^\alpha \operatorname{div} (u \otimes u) \quad \partial^\alpha D + \partial^\alpha \operatorname{curl} (u \otimes u) \partial^\alpha \Omega \right) \\ &\leq \frac{1}{4} (\|\partial^\alpha D\|^2 + \|\partial^\alpha \Omega\|^2) + C\|u \otimes u\|_{m+1}^2 \\ &\leq \frac{1}{4} (\|\partial^\alpha D\|^2 + \|\partial^\alpha \Omega\|^2) + (E_m^\varepsilon)^2. \end{aligned}$$

The second one is :

$$\begin{aligned} &\frac{1}{\varepsilon} \int \partial^\alpha \operatorname{div} (\rho u) \partial^\alpha \rho \\ &= \frac{1}{\varepsilon} \int u \cdot \nabla \partial^\alpha \rho \partial^\alpha \rho + [\partial^\alpha, u \cdot \nabla] \rho \partial^\alpha \rho + \partial^\alpha (\rho \operatorname{div} u) \partial^\alpha \rho \\ &= I + II + III. \end{aligned}$$

By an integration by parts, we have $I = -\frac{1}{\varepsilon} \int \operatorname{div} u |\partial^\alpha \rho|^2$, hence

$$I \leq \|\operatorname{div} u\|_{0, \infty} E_m^\varepsilon \leq (E_m^\varepsilon)^{\frac{3}{2}}.$$

The terms II and III are easily estimated by

$$\begin{aligned} II &\leq C\|u\|_m \frac{\|\rho\|_m^2}{\varepsilon} \leq C(E_m^\varepsilon)^{\frac{3}{2}}, \\ III &\leq C(\|u\|_m + \|\nabla u\|_m) \frac{\|\rho\|_m^2}{\varepsilon} \leq C(\|d\|_m + \|\omega\|_m) \frac{\|\rho\|_m^2}{\varepsilon} \leq C(E_m^\varepsilon)^{\frac{3}{2}}. \end{aligned}$$

Therefore we obtain the estimate

$$(16) \quad \frac{1}{\varepsilon} \int \partial^\alpha \operatorname{div} (\rho u) \partial^\alpha \rho \leq C(E_m^\varepsilon)^{\frac{3}{2}}.$$

To estimate the commutator, we have

$$\begin{aligned}
& \left(\Sigma^\varepsilon [\partial^\alpha, \mathcal{A}(x, \partial_x)] \mathcal{U}, \mathcal{U} \right) = \frac{1}{\varepsilon} \int [\partial^\alpha, u^{NS} \cdot \nabla] \rho \partial^\alpha \rho \, dx \\
(17) \quad & \leq C \frac{\|\rho\|_m}{\varepsilon} \left(\|u^{NS}\|_m \|\nabla \rho\|_{0,\infty} + \|\nabla u^{NS}\|_{0,\infty} \|\nabla \rho\|_{m-1} \right) \\
& \leq \mathcal{C} \left(\|u^{NS}\|_{1,\infty} + \|u^{NS}\|_m \right) E_m^\varepsilon.
\end{aligned}$$

To get the last line, we have used the fact that $m > 1 + d/2$.

Finally, we have that

$$\begin{aligned}
(18) \quad \varepsilon |(\Sigma^\varepsilon \partial^\alpha R, \partial^\alpha \mathcal{U})| & \leq \varepsilon \|\nabla u^{NS} \cdot \nabla u^{NS}\|_m \|d\|_m + \|\partial_t \operatorname{div} V^{NS}\|_m \|D\|_m \\
& \quad + \|\partial_t \operatorname{curl} V^{NS}\|_m \|\Omega\|_m \\
& \leq \|d\|_m^2 + \frac{1}{4} (\|D\|_m^2 + \|\Omega\|_m^2) + \varepsilon^2 \mathcal{C} (\|u^{NS}\|_{m+1} + \|\partial_t V^{NS}\|_m) \\
& \leq E_m^\varepsilon + \frac{1}{4} (\|D\|_m^2 + \|\Omega_m\|^2) + \varepsilon^2 \mathcal{C} (\|u^{NS}\|_{m+3}).
\end{aligned}$$

Notice that, to get the last line, we have used the explicit expression (2) of V^{NS} and the Navier-Stokes equations to replace the time derivative by spatial ones.

Now, we collect all the previous estimates (11)–(18) and we sum over α to find

$$\begin{aligned}
\frac{d}{dt} E_m^\varepsilon + \frac{1}{4} (\|D\|_m^2 + \|\Omega\|_m^2) & \leq \varepsilon^2 \mathcal{C} (\|u^{NS}\|_{m+3}) + \mathcal{C} (\|u^{NS}\|_{m+3}) E_m^\varepsilon \\
& \quad + (E_m^\varepsilon)^2 + C (E_m^\varepsilon)^{\frac{3}{2}} \\
& \leq \varepsilon^2 \mathcal{C} (\|u^{NS}\|_{m+3}) + \mathcal{C} (\|u^{NS}\|_{m+3}) E_m^\varepsilon + (E_m^\varepsilon)^2.
\end{aligned}$$

By using (9), we get with $M_\varepsilon \leq 1$ that

$$\frac{d}{dt} E_m^\varepsilon \leq \mathcal{C}_{m+3} \varepsilon^2 + \mathcal{C}_{m+3} E_m^\varepsilon, \quad \forall t \in [0, T^\varepsilon]$$

where $\mathcal{C}_{m+3} = \mathcal{C} (\|u^{NS}\|_{m+3})$ and hence by the Gronwall inequality we get that

$$(19) \quad E_m^\varepsilon(t) \leq \left(\alpha_m(\varepsilon) + t \mathcal{C}_{m+3} \varepsilon^2 \right) e^{\mathcal{C}_{m+3} t}, \quad \forall t \in [0, T^\varepsilon].$$

Consequently, if we choose $M_\varepsilon = (\alpha_m(\varepsilon) + T \mathcal{C}_{m+3} \varepsilon^2)^{\frac{1}{2}}$, we see that we cannot reach equality in (9) for $T^\varepsilon < T$. This proves that $T^\varepsilon \geq T$ and that (19) is valid on $[0, T]$.

REFERENCES

- [1] S. Alinhac and P. Gérard. *Opérateurs pseudo-différentiels et théorème de Nash-Moser*. Savoirs Actuels. [Current Scholarship]. InterEditions, Paris, 1991.
- [2] D. Aregba-Driollet, R. Natalini, and S.Q. Tang. Diffusive kinetic explicit schemes for nonlinear degenerate parabolic systems. *Math. Comp.* 73 (2004) 63-94.
- [3] M.K.Banda, A.Klar, L.Pareschi, M.Seaid. Compressible and Incompressible Limits for Hyperbolic Systems with Relaxation *Journal of Computational and Applied Mathematics*, 168 (2004) pp.41-52.
- [4] F. Bouchut and F. Golse and M. Pulvirenti. Kinetic equations and asymptotic theory. Series in Appl. Math., Gauthiers-Villars, 2000.
- [5] F. Bouchut, F. Guarguaglini, and R. Natalini. Discrete kinetic approximation to multidimensional parabolic equations. *Indiana Univ. Math. J.*, 49:723–749, 2000.
- [6] Y. Brenier, Convergence of the Vlasov-Poisson system to the incompressible Euler equations. *Comm. Partial Differential Equations* 25 (2000), no. 3-4, 737–754.
- [7] Y. Brenier, R. Natalini, and M. Puel. On a relaxation approximation of the incompressible Navier-Stokes equations. *Proc. Amer. Math. Soc.*, 132(4):1021–1028 (electronic), 2004.

- [8] S. Cordier and E. Grenier. Quasineutral limit of an Euler-Poisson system arising from plasma physics. *Comm. Partial Differential Equations*, 25(5-6):1099–1113, 2000.
- [9] D. Donatelli and P. Marcati. Convergence of singular limits for multi-D semilinear hyperbolic systems to parabolic systems. *Trans. Amer. Math. Soc.*, 356(5):2093–2121 (electronic), 2004.
- [10] D. Donatelli and P. Marcati. Diffusive singular limits and 3-D incompressible Navier-Stokes equation. To appear in *Proceedings of HYP 2004 Tenth International Conference on Hyperbolic Problems Theory, Numerics, Applications, Osaka, Japan*, Yokoama Publishers, Inc 2005.
- [11] E. Grenier. Defect measures of the Vlasov-Poisson system in the quasineutral regime. *Commun. Partial Differ. Equations* 20, No.7-8, 1189-1215, 1995.
- [12] E. Grenier. Oscillations in quasineutral plasmas. *Commun. Partial Differ. Equations* 21, No.3-4, 363-394, 1996.
- [13] Th. Katsaounis, Ch. Makridakis, C. Simeoni. Relaxation methods and finite element schemes for the incompressible Navier-Stokes equations. Preprint 2004.
- [14] G. Loeper. Quasineutral limit for the Euler-Poisson and Euler-Monge-Ampère systems. *Preprint*, 2003.
- [15] P. Marcati, A. Milani, and P. Secchi. Singular convergence of weak solutions for a quasilinear nonhomogeneous hyperbolic system. *Manuscripta Math.*, 60:49–69, 1988.
- [16] P. Marcati and B. Rubino. Hyperbolic to parabolic relaxation theory for quasilinear first order systems. *J. Differential Equations*, 162(2):359–399, 2000.
- [17] M. E. Taylor. *Partial differential equations. III*, volume 117 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1997. Nonlinear equations, Corrected reprint of the 1996 original.
- [18] Shu Wang. Quasineutral limit of Euler-Poisson system with and without viscosity. *Comm. Partial Differential Equations*, 29(3-4):419–456, 2004.